

# Electron Hydrogen scattering

2020-10-06

We use the Born approximation to compute the differential cross section for the elastic scattering of a fast electron by a hydrogen atom in the ground state. We will treat the hydrogen atom as a fixed target with a time-independent charge distribution.

blog: [Electron Hydrogen scattering](#) email: [quarktetra@gmail.com](mailto:quarktetra@gmail.com)

In Born approximation the differential cross section is given by

$$\frac{d\sigma}{d\Omega} = \left| \frac{m}{2\pi} \int d^3r e^{-i\mathbf{q}\cdot\mathbf{r}} V_I(r) \right|^2, \quad (1)$$

where  $V_I(r) = eV(r)$  is the interaction energy of the incoming electron with the hydrogen atom. It will be better to do the calculation in two parts since  $V(r)$  is contributed by two charge constituents, the proton and the electron. The contribution from the proton is straightforward to compute since we know  $V_p(r)$  right away:  $V_p(r) = \frac{e}{r}$ . The corresponding integral to be calculated as

$$I_1 \equiv \frac{m}{2\pi} \int d^3r e^{-i\mathbf{q}\cdot\mathbf{r}} \frac{e^2}{r}, \quad (2)$$

Unfortunately this integral is divergent, but we can regularize it by introducing an exponentially decaying function  $e^{-\epsilon r}$ , with  $\epsilon > 0$ . What we do is to evaluate the integral with  $V_p(r) = \frac{e^2 e^{-\epsilon r}}{r}$  and let  $\epsilon \rightarrow 0$  in the final answer (unlike physicists, mathematicians would strongly object: “You cannot change the order of integration and the limit  $\epsilon \rightarrow 0$ ”. We can think of it as assigning a finite number to a divergent integral by the renormalization procedure. Physically  $e^{-\epsilon r}/r$  terms represents the potential due a massive particle of mass  $\epsilon$ .) In this case we have

$$\begin{aligned} I_1 &\equiv \frac{me^2}{2\pi} \int d^3r e^{-i\mathbf{q}\cdot\mathbf{r}} \frac{e^{-\epsilon r}}{r} = 2me^2 \int dr \frac{\sin(qr)}{q} e^{-\epsilon r} = \frac{2me^2}{q} \Im \left\{ \int_0^\infty dr e^{-r(\epsilon - iq)} \right\} \\ &= \frac{2me^2}{q} \Im \left\{ \frac{1}{\epsilon - iq} \right\} = \frac{2me^2}{q^2}, \end{aligned} \quad (3)$$

where we finally set  $\epsilon = 0$ .

The second contribution comes from the electron cloud. The potential due to electron in the ground state is given as

$$V_e(r) = e\left(\frac{e^{-r/a_0}}{r} - \frac{1}{r} + \frac{1}{a_0}e^{-r/a_0}\right). \quad (4)$$

The first idea would be to plug this into the integral, and have fun... It can be done, but let's try something else,

$$I_2 \equiv \frac{me}{2\pi} \int d^3r e^{-i\mathbf{q} \cdot \mathbf{r}} V_e(r) = \frac{2me}{q} \int_0^\infty r V_e(r) \sin(qr) dr \quad (5)$$

Now the idea is the integration by parts twice with the definitions  $\mathcal{U} = rV(r)$  and  $d\mathcal{V} = \sin(qr)$  for the first one and  $\mathcal{U} = \frac{d}{dr}(rV(r))$  and  $d\mathcal{V} = \cos(qr)$  for the second one. Note that we can drop  $\mathcal{UV}|_0^\infty$  since  $\mathcal{U}$  vanishes at the boundaries. After two integration by parts we get

$$I_2 = -\frac{2m}{q^3} \int_0^\infty \frac{d^2}{dr^2}(rV_e(r)) \sin(qr) dr. \quad (6)$$

What we have achieved is  $\frac{d^2}{dr^2}(rV_e(r))$  term, which is  $r \nabla^2 V_e$ . By Poisson's equation it can be replaced by  $r(-4\pi\rho_e(r))$  where

$$\rho_e(r) = -e|\psi(r)|^2 = -\frac{e}{\pi a_0^3} e^{-2r/a_0} \quad (7)$$

So the integral we need to deal with is

$$\begin{aligned} I_2 &= -\frac{8me^2}{a_0^3 q^3} \int_0^\infty r e^{-2r/a_0} \sin(qr) dr = -\frac{8me^2}{a_0^3 q^3} \Im \left\{ \int_0^\infty r e^{-(2/a_0 - iq)r} dr \right\} \\ &= -\frac{8me^2}{a_0^3 q^3} \Im \left\{ \left( \frac{-d}{d\alpha} \right)_{\alpha=(2/a_0 - iq)} \int_0^\infty e^{-\alpha r} dr \right\} \\ &= -\frac{8me^2}{a_0^3 q^3} \Im \left\{ \frac{qa_0}{(2 - iqa_0)^2} \right\} = -\frac{32me^2}{[4 + (qa_0)^2]^2 q^2}. \end{aligned} \quad (8)$$

Combining both terms we have

$$\frac{d\sigma}{d\Omega} = \frac{4m^2 e^4}{q^4} \left( 1 - \frac{16}{[4 + (qa_0)^2]^2} \right)^2. \quad (9)$$

Before we discuss the limiting cases, let's solve the problem one more time using a shortcut. The object we are dealing with is,

$$I = \int d^3r e^{-i\mathbf{q} \cdot \mathbf{r}} V(r), \quad (10)$$

which is nothing but the Fourier transform of the potential energy. Instead of jumping onto the problem head on, let's follow a detour. Consider the Poisson's equation,

$$\nabla^2 V(r) = -4\pi\rho(r) = -4\pi e(\delta(\vec{r}) - \rho_e(r)). \quad (11)$$

Instead of Fourier transforming the potential itself, we can Fourier transform Eq. 11 which gives

$$-q^2\tilde{V}(q) = -4\pi e(1 - F\{\rho_e(r)\}) \rightarrow \tilde{V}(q) = \frac{4\pi e(1 - F\{\rho_e(r)\})}{q^2}. \quad (12)$$

$F\{\rho_e(r)\}$  term is to be calculated by usual means, and it reproduces the second term in Eq 9.

For large  $qa_0$  the second term in Eq. 9, which accounts for the electron cloud, can be neglected. So the incoming electrons are deflected only by the proton. For small  $qa_0$  the cross section becomes zero which is a manifestation of the fact that the hydrogen atom is neutral. This means that only highly deflected electrons can probe into the electron cloud, and “see” the proton.